

## Pelvic Floor Physical Therapy in the Treatment of a Patient With Interstitial Cystitis, Dyspareunia, and Low Back Pain: A Case Report

Nicole Cozean, PT, DPT, WCS, CSCS

### ABSTRACT

**Background:** Patients with a diagnosis of interstitial cystitis (IC) often have other related symptoms and conditions caused by a hypertonic pelvic floor. This case report examines the physical therapy diagnosis, management, and outcomes of a patient with IC, dyspareunia, and low back pain.

**Study Design:** A single-subject case report on the initial examination, interventions, and outcomes of the pelvic floor physical therapy for a patient with a common triad of pelvic floor conditions. The case report includes a detailed description of each session and patient response to guide interventions in similar cases.

**Case Description:** This case follows the management of a patient with a diagnosis of IC accompanied by urgency and frequency of urination, severe dyspareunia, and low back pain. Interventions included a combination of orthopedic and pelvic floor physical therapy, internal and external manual therapy, behavioral therapy, therapeutic exercises, and education.

**Outcomes:** After 19 physical therapy visits over 3 months, the patient was able to resume full-time employment and sexual intercourse with little to no pain. Pain was reduced by nearly 80% on the Short-Form McGill Pain Questionnaire. On evaluation, the patient scored below average on 7 SF-36 quality-of-life subcategories, which improved to above-average scores in 7 of the 8 subcategories at discharge.

**Discussion:** The diagnosis of IC is often accompanied by other associated pelvic floor conditions. A combination of pelvic floor physical therapy and orthopedic techniques may be necessary to relieve urinary frequency and urgency, nocturia, bladder pain, dyspareunia, and low back pain in complex patients with IC.

**Key Words:** dyspareunia, interstitial cystitis, low back pain, manual physical therapy, painful intercourse, pelvic floor physical therapy

### BACKGROUND

Many women with interstitial cystitis (IC) have coexisting pelvic dysfunction and symptoms, includ-

ing pelvic floor muscle (PFM) spasm, low back pain (LBP), and dyspareunia, complicating physical therapy management and outcomes. Characterized by frequency and urgency of urination, dysuria, and chronic pain of the bladder, IC remains a disease diagnosed primarily by exclusion.<sup>1,2</sup> Typically, urinary tract infection and bladder cancer are ruled out before the patient receives a diagnosis of IC. As one review states, “A broad clinical definition of IC includes any patient who complains of urinary urgency, frequency, and/or pelvic/perineal pain, in the absence of any identifiable cause, such as bacterial infection or carcinoma.”<sup>2(p328)</sup>

It has recently been determined that the prevalence of IC is much higher than was previously thought. A survey of nearly 150 000 households in the United States found that between 3.3 million and 7.9 million people in the United States are afflicted with IC.<sup>1</sup> The diagnosis of IC is frequently linked to a diagnosis of sexual pain or dysfunction<sup>2</sup>; 90% of women with a diagnosis of IC had experienced 1 or more incidences of sexual dysfunction in the previous 4 weeks.<sup>3</sup>

It is well established that PFM spasm is present in patients with irritable voiding diagnoses such as IC in addition to other pain diagnoses such as dyspareunia, LBP, and sacroiliac joint (SIJ) dysfunction.<sup>4-15</sup> Frequently, the PFMs are overactive in an attempt to stabilize the painful and dysfunctional lumbar spine, hip, and/or pelvic girdle.<sup>12,16-21</sup> PFMs with trigger point presence can refer pain to other pelvic girdle structures and can also mimic low back and coccyx pain.<sup>16-18</sup>

Typical treatments of IC include analgesic installations into the bladder, oral analgesics, behavioral therapy, and pelvic floor physical therapy.<sup>2,22,23</sup> Bladder installations have been demonstrated to be helpful for some patients, but others report a worsening of symptoms following the invasive procedure and catheterization. Currently, an installation of dimethyl sulfoxide

*PelvicSanity, Laguna Hills, California.*

*The author declares no conflicts of interest.*

DOI: 10.1097/JWH.0000000000000062

is the only Food and Drug Administration–approved installation, but heparin, lidocaine, or liquid pentosan-polysulfate (Elmiron) are also frequently used.<sup>24</sup> A placebo-controlled study showed that only 40% of patients were helped with intravesical Elmiron as compared with 20% for placebo.<sup>25</sup> Alkalinized lidocaine showed an improvement in 30% of patients after 8 days of treatment as compared with 9% in the placebo group.<sup>26</sup> A combination of heparin and lidocaine was more promising, showing symptom relief in 75% of patients, but no long-term follow-up was conducted.<sup>27</sup> Prescription oral medication typically consists of Elmiron, which not only has shown small but statistically significant improvements in urgency, frequency, and pain with urination but also has significant side effects and can take up to 6 months of treatment before the patient sees therapeutic effect.<sup>28</sup> Patients also use over-the-counter analgesics, typically nonsteroidal anti-inflammatory drugs, in an attempt to reduce pain associated with the condition. Muscle-relaxant suppositories such as baclofen and diazepam are routinely used for PFM spasm; when combined with pelvic floor physical therapy, these suppositories have been shown to decrease pain and muscle spasm in patients with pelvic floor dysfunction and urogenital pain disorders.<sup>29</sup>

Pelvic floor physical therapy has been demonstrated, alone or in conjunction with other methods of treatment, to be effective in reducing the symptoms associated with IC and PFM spasm.<sup>8,11,20,30–32</sup> In a 2001 study involving 10 patients with a diagnosis of IC, Weiss<sup>33</sup> showed that 70% (7 of 10) had moderate (>50%) or marked (>75%) improvement with manual myofascial release techniques. Another 2001 study demonstrated that 90% of patients with IC saw benefits from transvaginal Thiele massage.<sup>34</sup> A study showed that 57% of the patients who received myofascial pelvic floor physical therapy demonstrated moderate or marked improvement as compared with 21% of those in the control group.<sup>30</sup>

While these reports demonstrate the benefit of pelvic floor physical therapy for the traditional symptoms of IC—pelvic pain and urinary symptoms—they do not address whether the improvements extend to associated symptoms. This case report focuses on the pelvic floor physical therapy management of a woman with IC, PFM spasm, and LBP—a common set of coexisting conditions in patients with pelvic floor disorders. Clinical trials evaluating pelvic floor physical therapy also present data from a discrete end point of the trial. This case report tracks the progress of the patient through each visit of her physical therapy regimen, demonstrating reductions in pain levels, improvements in urinary symptoms, restoration of the activities of daily living, and improvements in quality of life.

## CASE DESCRIPTION (HISTORY, SYSTEMS REVIEW, TESTS, AND MEASURES)

### Patient History

The patient, a 43-year-old woman (G2P2), was referred to physical therapy from an urogynecologist. The patient history was taken in an oral interview format, guided by the Pelvic Pain Assessment form created by the International Pelvic Pain Society.<sup>35</sup> The patient reported severe urgency and frequency of urination with burning urethral pain, as well as a “dull ache” in her lower back and suprapubic region coinciding with a full bladder that significantly affected daily life. In addition, she complained of deep pain with sexual intercourse lasting up to days after, and urinary urgency during, intercourse. Nocturia was reported, with the patient waking 5 to 6 times each night. The patient reported the onset of her symptoms approximately 6 months prior to the initial physical therapy evaluation, with progressive worsening of pain and symptoms forcing her to seek medical care. When further questioned about the specifics of symptom onset, the patient reported similar symptoms with decreased intensity and frequency (both urologic and back and pelvic pain) since giving birth more than 15 years ago. Vaginal delivery has shown to cause partial denervation of the pelvic floor (with consequent reinnervation) in most primiparous women, which may pave the way for pelvic floor dysfunctions later in life.<sup>36</sup> During the examination, the patient also complained of severe back pain that had increased since the onset of her voiding symptoms, disturbing her sleep and causing severe pain with sitting and any form of supine or side-lying positions.

Aggravating factors to her pain included sexual intercourse, vaginal and labial contact with clothing, full bladder, stress, urination, sitting, and standing. Easing factors to her pain included bowel movement, stool softeners, heating pad, and hot bath. The patient had a history of constipation, which has been shown to exacerbate bladder and back pain.<sup>37,38</sup>

The patient had participated in yoga or other exercise 5 or 6 times per week, although her activities had been curtailed because of the increase in her symptoms, particularly her LBP. With a pending IC diagnosis, she reported following an IC diet,<sup>39</sup> eliminating caffeine, alcohol, and other trigger foods. She also reported limiting water intake due to urinary urgency and frequency.

In addition, functional deficits and negative effects on social life and coping mechanisms were addressed in the patient interview, as these are a common by-product of this triad of diagnoses.<sup>4,5</sup> Prior to symptom presentation, the patient was a full-time social worker. Upon initial evaluation, she was unable to work and had taken a medical leave of absence due to the severity

of her symptoms. She reported that her relationship with her husband was strained not only because of her inability to have sexual intercourse but also with her recent role change in the household (going from full-time employment to not working). She was having significant difficulty caring for her family, sleeping, and performing other aspects of daily life due to severe pain as well as severe urinary urgency and frequency. She self-reported depression over onset of symptoms, and she also reported a supportive group of family and friends in addition to seeking help from a psychologist when needed during these challenging times.

### Medications

The patient had seen 4 physicians, all of whom had recommended or prescribed different medications for pain and symptom relief. The patient was being treated with bladder instillations of heparin and bupiracaine 1 or 2 times per week by the referring urogynecologist, with only slight relief experienced. She was prescribed acetaminophen/hydrocodone by her primary care physician for the LBP, but the patient reported using the medication sparingly due to the side effects of nausea and constipation. Zolpidem had also been prescribed by another physician for pain-related insomnia. Her gastroenterologist had prescribed polyethylene glycol for constipation, which had not eliminated this symptom. The patient had also started baclofen vaginal suppositories 1 day prior to evaluation, prescribed by her referring urogynecologist. The patient chose to not try Elmiron at this time.<sup>28</sup>

### Medical/Surgical History/Systems Review

Pertinent medical history included a history of bulimia and laxative abuse, for which she had sought treatment. She indicated control over her eating disorder for more than 20 years at the time of initial evaluation. She reported a 10-year history of intermittent lower back pain and dysmenorrhea. The patient had undergone a breast-reduction surgery approximately a year ago and a loop electrosurgical excision procedure more than 10 years prior to the initial evaluation. She had 2 vaginal deliveries, each requiring an episiotomy. No red flags were present in the patient's systems review, and vital signs were within normal limits.

## TESTS AND MEASURES

### Outcome Measures

The SF-36 was administered to measure quality of life.<sup>40</sup> The SF-36 includes 8 subcategories or subscales, where a category score below 50 is considered below average. The Short-Form McGill Pain Questionnaire was also administered to the patient.<sup>41</sup> Responses on the Short-Form McGill Pain Questionnaire outcomes

are rated on a 4-point scale (none = 0; mild = 1; moderate = 2; severe = 3) for 15 different descriptors. The first 11 describe sensations, whereas the final 4 describe patient feelings, with a possible maximum severe pain rating being 45. The patient's pain was also assessed on the Verbal Rating Scale Pain Scale (1-10 scale).

### External Assessment

The patient gave informed consent before initiating examination, including consent to examine external and internal pelvic structures.<sup>42</sup> Her posture was not one typically associated with pelvic pain (characterized by excessive anterior pelvic tilt, lumbar lordosis, and thoracic kyphosis)<sup>10</sup> but did represent consistency with LBP classification of being extension-sensitive.<sup>43</sup> Regardless, a finding of faulty posture is consistent with the literature. Women with chronic pelvic pain have faulty posture, movement patterns, and body awareness compared with women without pelvic pain.<sup>43</sup>

Her posture faults included forward head and rounded shoulder posture, with a posterior pelvic tilt in the standing and sitting positions. She showed overactive bilateral gluteal recruitment in standing. In standing, her left iliac crest and gluteal fold were elevated compared with the right. Supine bony landmark palpation revealed left ilial posterior rotation with outflare, elevated left pubic symphysis height, and tenderness with palpation along left suprapubic ramus. SIJ and postural assessment was performed per Richard Jackson's and Diane Lee's examination techniques to assess the potential effects of faulty biomechanics of the pelvis affecting the pelvic floor.<sup>21,44-46</sup> The patient was found to have functional SIJ dysfunction. She demonstrated left SIJ hypomobility along with the positional fault of the ilium on sacrum. Gross assessment of trunk range of motion in flexion/extension, side-bending, and lateral rotation was all found to be within normal limits.<sup>47,48</sup> The patient experienced pain at end range of lumbar extension and was exhibiting pain avoidance postures to bias toward lumbar flexion.<sup>49</sup>

Bilateral hamstring length, measured by the 90-90 hamstring test, revealed bilateral limitations ( $-45^{\circ}$  R;  $-30^{\circ}$  L).<sup>47</sup> All other lower extremity range of motion was within normal limits<sup>47,50</sup> except hip external rotation, which was measured at  $40^{\circ}$ , whereas hip internal rotation was measured at  $35^{\circ}$  bilaterally. In the active straight leg raise (ASLR) test, the patient showed increased ability to perform ASLR on the left with anterior compression of pubic symphysis, indicating transversus abdominis weakness.<sup>51</sup>

The patient demonstrated hypomobility of L2 and L3 via posterior-to-anterior spring testing, hypermobility of L4, capsular end feel of L2-L3, and painful end feel of L4.<sup>48</sup> Marcher's test showed a positive left

SIJ ilial on sacral motion with a positive Trendelenburg sign in single-limb stance.<sup>44</sup> In the straight leg raise test for sciatic nerve tension testing, the patient presented with negative results for crossover sign, indicating no lumbar disc pathology.<sup>50</sup> Passive straight leg raise test was positive at 35° for sciatic nerve irritation, with primary mechanical interface of the nerve at the proximal hamstring and piriformis.<sup>50</sup>

Lower extremity strength,<sup>52</sup> upper extremity strength,<sup>45,47,50</sup> and upper extremity range of motion<sup>47</sup> were within normal limits. Test of diastasis recti abdominis<sup>21</sup> was negative. The patient's deep tendon reflexes were also found to be within normal limits, lower extremity sensation was intact to light touch and sharp/dull, and perineal sensation was intact to light touch.

### Pelvic Floor Internal Examination

A thorough pelvic floor internal examination consists of an assessment of PFM strength, presence of trigger points, and symptom reproduction to palpation of the pelvic floor musculature. The quantification of a pelvic floor contraction is problematic especially in patients with PFM spasm. There is no validated scale to quantify contractions of the PFMs in patients with a hypertonic pelvic floor; therefore, quantification—more than absent, weak, normal, or strong—is not recommended.<sup>53</sup> During assessment of PFM hypertonicity and symptom reproduction, a severe or strong rating was given when pain associated with palpation met or exceeded 7/10 on the self-reported pain scale. Stiffness in the musculature and the presence of any twitch, spasm, or trigger point were also noted. Often resting tone is measured with a transvaginal electromyography assessment.<sup>54</sup> But the patient had significant pain, thus electromyography was not used to prevent exacerbation of symptoms.

PFM strength<sup>53</sup> was strong, with a muscle endurance hold of 5 seconds. However, she had difficult relaxing PFMs after contraction, indicating poor PFM coordination. Severe PFM trigger points and pain were located in the left of superficial transverse perineal, PC (pubococcygeus), IC (iliococcygeus), OI (obturator internus), piriformis and coccygeus, and urogenital diaphragm (UGD) muscles. Urinary urgency was reproduced with palpation of bilateral IC-OI interface at ischial spine and bilateral UGD muscles. Pelvic floor hypertonicity was found to be moderate (pain associated with palpation met 4-6/10 of the pain scale) on the right superficial transverse perineal, PC, IC, OI, piriformis and coccygeus, and UGD muscles. Palpation of bilateral deep IC, OI, and coccygeus reproduced (left greater than right) pain similar to that during intercourse. She reported increased tenderness of pelvic clock at introitus from 2 to 6 o'clock but demonstrated a negative Q-Tip test.<sup>55,56</sup>

Myofascial assessment for trigger points of the pelvic girdle muscles revealed pain and trigger point presence with palpation of the bilateral quadratus lumborum, lumbosacral multifidus, bilateral piriformis, bilateral gluteus maximus and medius, and bilateral iliopsoas, bilateral short and long adductors, and proximal hamstrings. The assessment revealed significant restriction of ischiorectal, suprapubic, and abdominal fascia, with increased tissue tension of sacrospinous and sacrotuberous ligaments.<sup>8</sup>

## EVALUATION AND PHYSICAL THERAPY DIAGNOSIS

The patient presented with moderate to severe pelvic floor tension myalgia/muscle spasm that contributed to urinary urgency and frequency, pain with intercourse, and suprapubic pelvic pain. Symptoms and PFM spasm were complicated by left SIJ dysfunction, a history of LBP, and current LBP flare, in addition to core musculature and hip muscle weakness. During examination, most of the patient's IC/irritable voiding symptoms were reproduced with PFM palpation, indicating PFM spasm as the dominant pain mechanism in this case.

### Prognosis

Because of the patient's relatively recent worsening of symptoms, prognosis was deemed good to excellent for full resolution of all pelvic pain symptoms and LBP. Because of the chronic nature of IC, the patient will likely always be susceptible to irritable voiding symptoms but would likely see a significant decrease in urinary urgency and frequency symptoms since these symptoms were reproduced with PFM palpation and presence of PFMS.

## PHYSICAL THERAPY INTERVENTION

Manual therapy was the primary physical therapy modality used and consisted of manual trigger point release, myofascial release, and Thiele massage to transvaginal PFMs.<sup>32</sup> Other techniques used for surrounding areas at the pelvic girdle were myofascial release (including skin rolling), connective tissue mobilization, joint mobilization (SIJ, hip, lumbar, and thoracic spine as indicated), neural glides (for both sciatic and pudendal nerves), muscle energy techniques, contract/relax techniques, and visceral mobilization.\* These interventions were chosen to release PFM spasm and trigger points, as noted in the initial examination, as well as addressing dysfunction from the pelvic girdle to attend to LBP.

\*References 2,6-8,14,16,17,20,21,50.

Therapeutic exercises were selected on the basis of the patient's needs in lengthening and strengthening found on the evaluation. Therapeutic exercises began with pelvic floor relaxation stretches and diaphragmatic breathing instruction. The first stretch taught to the patient was the "Happy Baby" yoga pose, which both relaxes the pelvic floor and biases toward lumbar flexion. The patient was progressed to a deep squat stretch when symptoms and hip muscle length allowed. As the patient's pelvic floor musculature began to respond to manual therapy and was less prone to painful flares, she was progressed to core stabilization exercises, including transverse abdominis and lumbosacral multifidus recruitment, as well as incorporating other pelvic girdle structures and hip stabilizers into exercises. The patient was monitored throughout sessions and was progressed accordingly based on strength and flexibility improvements and response to exercise throughout the program. The patient's exercise position was modified for a flexion bias, as she was found to be positionally sensitive to lumbar extension on initial evaluation.<sup>49</sup> The importance of rehabilitating multifidus and transversus abdominis muscles in patients with LBP is well documented.<sup>57,58</sup>

Diaphragmatic breathing was implemented as an exercise unto itself in addition to being incorporated into core stabilization. Chaitow<sup>59,60</sup> points out this importance in an article by Key et al that cautions the therapist to be aware of "core rigidity" when core stabilization exercises are taught to the patient without functional movement patterns and proper breathing. This patient was monitored frequently for proper breathing patterns during therapeutic exercise in addition to using diaphragmatic breathing as a pelvic floor relaxation exercise. There is a clear connection between respiratory function, pelvic floor function and tone, and SIJ stability, particularly in women.<sup>61</sup> Examination of this patient demonstrated objective evidence that all 3 of these components needed to be addressed in her exercise program.

The patient was educated with visual aids in the anatomy and physiology of the pelvic girdle and PFMs, and how symptom presentation may have neuromuscular and musculoskeletal contributions. She was also informed about proper water intake,<sup>62</sup> keeping a voiding diary,<sup>63</sup> urge suppression training techniques, proper body mechanics, and home exercise program. Home exercises included colon massage to relieve constipation,<sup>38</sup> crystal wand instruction for self-treatment of transvaginal PFM trigger point release,<sup>64</sup> pelvic girdle-associated muscle stretching, and core stabilization program modified for patient's yoga classes.

## OUTCOMES

The patient reported and maintained consistent decreases in LBP, urinary urgency and frequency, and PFMS, while demonstrated an increase in function and return to normal activities. The patient had 1 flare due to admittedly abandoning home program and IC diet over family-based weekend. The patient was able to identify reasons for flare and demonstrate self-correction techniques to decrease symptoms (Table 1).

Upon discharge, patient reported substantial improvement in all symptoms. Urinary urgency and frequency were reduced to normal levels, and urethral pain with urination was also greatly reduced. Low back and SIJ pain was completely eliminated, and suprapubic pain was substantially reduced upon discharge. Vaginal pain with intercourse and postintercourse was reported to be eliminated or only mildly bothersome according to the patient (Table 2).

With the improvement in her condition, the patient was able to sit and stand for more than 2 hours, which has allowed her to resume many activities of daily living, including outings with her family and returning to her work as a social worker. She resumed sexual intercourse with her husband, with no discomfort or occasionally mild, intermittent pain. Nocturia was reduced to twice per night, improving the patient's sleep. She was able to return to her regular exercise routine of yoga 5 times a week. The patient reported only mild and intermittent symptoms and was pleased with her progress. She was confident her improvement would continue with consistency of home exercise program and self-treatment techniques (Figures 1 and 2).

This patient demonstrated significant improvement on the SF-36 quality-of-life measurement. As per the *User's Manual for the SF-36(v2) Health Survey*, norm-based scoring should be used to simplify interpretation. The general population norms are built into the scoring algorithm, which makes comparing this patient with normative scores valuable. In this case, the patient scored below average on 7 of 8 categories upon initial evaluation and above average on 7 of 8 categories upon discharge from physical therapy (see Figure 1).

She also demonstrated a decrease on Short-Form McGill Pain Questionnaire, indicating a substantial improvement in pain symptoms. Her overall pain score was reduced from 29 of 45 to 6 of 45 at discharge. Substantial improvements were seen in both sensory (4-fold reduction) and affective (8-fold reduction) categories.

**Table 1. Summary of Treatments and Visit Outcomes**

Visit	Manual Therapy	Education	Therapy Ex	Patient Reported Progress and Visit Notes
1	X	X	X	(Evaluation)
2	X	X		Patient used muscle-relaxant suppository prescribed by the physician; reported 50% reduction in pain by the end of the session and able to achieve comfortable position in supine.
3	X	X	X	Patient reported a continued 50% reduction in back pain; diagnosed with bladder infection; therefore, internal manual therapy held.
4	X	X	X	External treatment performed because of clearing bladder infection; patient reported continued decrease in back pain with improved sleep and urgency at night.
5	X	X	X	Patient continue >50% LBP improvement, and no longer reported LBP as her primary symptom; within-session improvement in urgency reproduction with internal PFM palpation from 8/10 (intensity) to 2-3/10.
6	X	X	X	Patient reported improved sleep due to decreased pain, LBP 85% gone; urinary urgency persists despite within-session improvements of urgency reproduction and decreased PFM spasm. Cystoscopy results: 75% bladder lining inflamed consistent with bladder irritation and IC.
7	X	X	X	Patient reported no LBP, and first decrease in urinary urgency.
8	X	X	X	During the session, trigger point reproduction of urinary urgency decreased from 5/10 to 1/10 with palpation of PFMs.
9	X	X		Patient reported overall improvement with urgency, now rating at 3-4/10 (vs 8/10 on initial evaluation). The patient was able to return to yoga exercise now that LBP was eliminated as a symptom.
10	X	X		Urgency decreased from 4/10 to 1-2/10 within session. The patient is continuing with yoga and core strengthening exercises. Worked to modify diet and home program to remove some restrictions to fit her current status.
11	X	X		Urgency decreased to 2/10; able to continue to increase yoga exercises.
12	X	X		Functional improvements continued, and the patient started to look for a new job and increased yoga to 3× per week. Sustained improvements in urgency even while “cheating” on diet.
13	X	X		Urgency improved, even with normal food triggers; job interview completed; urgency reduced to 2/10 with normalization of frequency.
14	X	X	X	Maximum intensity of urinary urgency reported as 1-2/10; patient began part-time work.
15	X	X		Symptoms consistently mild; the patient was confident in her return to work. Urgency no longer reproduced with PFM palpation.
16	X	X	X	Flare reported with increased urgency/frequency of urination after being “bad with diet over whole weekend,” no yoga or home exercises.
17	X	X	X	Reduced urination to 1× per 1.5-2 h, 3-4× at night after the recent increase due to flare.
18	X	X	X	Mild to WNL PFM spasm; progressed to full-time work now; only mild intermittent symptoms.
19	X	X	X	Full-time work; discharged, 0-3/10 symptoms; intermittent mild urgency controlled by home exercise and self-treatment.

Abbreviations: LBP, low back pain; pelvic floor muscle; WNL, within normal limit.

**DISCUSSION**

At the time of her initial physical therapy evaluation, it was clear that the patient’s primary symptoms, pain, and functional limitations were related to her acute LBP. Concurrently, her PFMs were extremely hypersensitive, likely due to not being able to lie supine without pain. For those reasons, the initial

focus was on relieving the acute pain to allow for proper positioning and reducing the internal hypersensitivity of tissues. Manual therapy for the first 4 visits consisted of correction of pelvic obliquity, SIJ mobilization, neural mobilization of the sciatic nerve, and lumbar spine joint mobilizations at grades I and II for inflammation reduction. Manual therapy during these visits was externally focused and consisted

**Table 2. Improvements in Pain and Function**

	Pretreatment	At Discharge
Urgency and frequency of urination	>16 times per day	<8 times per day
Low back and suprapubic pain	9/10	2/10
Burning urethral pain with urination	2/10	1/10
Burning vaginal pain introitus sexual intercourse	8/10	0-3/10
Deep vaginal pain with intercourse	9/10	0-3/10
Pelvic pain last 3 d after intercourse	7/10	0-2/10
Low back/SIJ pain	10/10	0/10
Pain in groin and suprapubic region when lifting	6/10	2/10
Abbreviation: SIJ, sacroiliac joint.		


of tissue approximation techniques, rather than internal trigger point release, which was too painful for the patient to tolerate at that time. After 5 visits, the patient reported substantial reductions in LBP.


After the patient's LBP had been reduced, urinary frequency and urgency became the patient's primary symptoms, and the pelvic floor hypersensitivity abated to the point where internal therapy was tolerable for the patient. Trigger points within the pelvic floor, primarily the urogenital diaphragm muscles, directly reproduced the patient's specific urinary urgency. Sexual pain was reproduced with palpation of levator ani muscles. Visits 5 to 15 focused on manual therapy to decrease pelvic floor trigger points, which resulted in a decrease in urinary symptoms and sexual pain.

This intervention was maintained because the patient continued to report improvements, both within and between sessions, even as the patient resumed work, exercise, and sexual intercourse. As expected, as she increased her activities of daily living, there was a corresponding increase in PFM trigger point presence, which was then addressed in treatment to allow her to continue to progress her activity level in a systematic fashion.

In the next stage of management, from visits 16 to 19, sessions focused on eradicating any residual and even mild SIJ dysfunction or LBP, as well as any persisting PFM trigger points, while the patient continued to progress toward full prior level of function, including full-time employment without symptoms. Visit frequency was reduced during this time. Additional focus was placed on core stabilization and educating the patient on independent management of flares and symptoms. The patient was educated on the practical implications of her long-term condition, including changes in diet and exercise. She was taught how to assess and respond to flares and increases in sexual pain, including modifying sexual positioning to minimize symptoms. Additional education included maintenance and self-interpretation of a voiding diary, trigger point release using the crystal wand, and pelvic floor relaxation.

Upon discharge following visit 19, the patient had returned to full-time employment at time of discharge, with no exacerbation of symptoms. She returned to her regular exercise program of high-intensity Bikram yoga 5 times per week. She was able to engage in regular, non- or mildly painful sexual intercourse with her husband. Her back pain and SIJ pain were completely eliminated, with occasional

**Figure 1.** Improvement in SF-36 quality-of-life questionnaire.



**Figure 2.** Reduction in pain on Short-Form McGill Pain Questionnaire.

mild pain. Overall, she reported a significant increase in confidence in being able to independently manage any residual symptoms.

Additional research is needed in this patient population. The prevalence of this symptom presentation—LBP, sexual pain, and urinary urgency and frequency, and bladder pain—has not been well established. It is unclear whether there are any casual or risk factors in the relationship between orthopedic dysfunction in the pelvic girdle and pelvic floor dysfunction. Interventions for this complex symptom presentation are still subjective, and a standardized treatment protocol has not yet been developed. Further large-scale studies on pelvic floor physical therapy intervention are needed.

## CONCLUSIONS

This case illustrates the benefits of a combination of orthopedic and pelvic floor physical therapy interventions to substantially improve outcomes for complex IC patients. Orthopedic and pelvic floor dysfunctions often coexist. Both must be addressed to alleviate the acute symptoms and allow for sustained improvement and return to full prior level of function. Appropriate and individualized physical therapy can result in rapid and sustainable improvement in pelvic and LBP, as well as urgency and frequency of urination, and lead to the patient’s resumption of the activities of daily living and ultimately improve her quality of life.

## REFERENCES

- Berry SH, Elliott MN, Suttorp M, et al. Prevalence of symptoms of bladder pain syndrome/interstitial cystitis among adult females in the United States. *J Urol.* 2011;186(2):540–544.
- Chaitow L. Chronic pelvic pain: pelvic floor problems, sacroiliac dysfunction and the trigger point connection. *J Bodywork Mov Ther.* 2007;11:327–339.

- Bogart LM, Suttorp MJ, Elliott MN, Clemens JQ, Berry SH. Prevalence and correlates of sexual dysfunction among women with bladder pain syndrome/interstitial cystitis. *Urology.* 2011;77(3):576–580.
- Bernstein A, Philips H, Linden W, et al. A psychophysiological evaluation of female urethral syndrome: evidence for a muscular abnormality. *J Behav Med.* 1992;15:299–312.
- Fall M, Baranowski A, Fowler C, et al. EAU guidelines for chronic pelvic pain. *Eur Urol.* 2004;46:681–689.
- Weiss JM, Prendergast SA. Screening for musculoskeletal causes of pelvic pain. *Clin Obstet Gynaecol.* 2003;46(4):773–782.
- Weiss JM. Chronic pelvic pain and myofascial trigger points. *Pain Clin.* 2000;2(6):13–18.
- Fitzgerald MP, Kotarinos R. Rehabilitation of the short pelvic floor, part II: treatment of the patient with the short pelvic floor. *Int Urogynecol J Pelvic Floor Dysfunct.* 2003;14(4):269–275.
- Baker PK. Musculoskeletal origins of chronic pelvic pain: diagnosis and treatment. *Obstet Gynecol Clin North Am.* 1993;20:719–742.
- King PM, Myers CA, Ling FW, Rosenthal RH. Musculoskeletal factors in chronic pelvic pain. *J Psychom Obstet Gynaecol.* 1991;12:87–98.
- Doggeweiler-Wygul R, Wiygul JP. Interstitial cystitis, pelvic pain, and the relationship to myofascial pain and dysfunction: a report on four patients. *World J Urol.* 2002;20:310–314.
- Sarton J. Physical therapy for pelvic pain: understanding the musculoskeletal connection. *Female Patient.* 2007;32:50–56.
- Sinaki M. Tension myalgia of the pelvic floor. *Mayo Clin Proc.* 1977;52:717–722.
- Bassaly R, Tidwell N, Bertolino S, Hoyte L, Downes K, Hart S. Myofascial pain and pelvic floor dysfunction in patients with interstitial cystitis. *Int Urogynecol J.* 2011;22(4):413–418.
- Hodges P, Richardson C. Inefficient muscular stabilization of the lumbar spine associated with low back pain. A motor control evaluation of transversus abdominis. *Spine.* 1996;21(22):2640–2650.
- Vleeming A, Stoockart R, Volkers ACW, Snijders CJ. Relation between form and function in the sacroiliac joint, part 1: clinical anatomical aspects. *Spine.* 1990;15(2):130–132.
- Vleeming A, Volkers ACW, Snijders C J, Stoockart R. Relation between form and function in the sacroiliac joint, part 2: biomechanical aspects. *Spine.* 1990;15(2):133–136.
- Sapsford R. Rehabilitation of pelvic floor muscles utilizing trunk stabilization. *Man Ther.* 2004;9(1):3–12.
- Sapsford RR, Hodges PW, Richardson CA, Cooper DH, Markwell SJ, Jull GA. Co-activation of the abdominal and pelvic floor muscles during voluntary exercises. *NeuroUrol Urodyn.* 2001;20:31–42.
- Lukban JC, Whitmore K, Kellogg-Spadt S, et al. The effect of manual physical therapy in patients diagnosed with interstitial cystitis, high-tone pelvic floor dysfunction, and sacroiliac dysfunction. *Urology.* 2001;57(6)(suppl):121–122.
- Lee D. *The Pelvic Girdle: An Approach to the Examination and Treatment of the Lumbo-Pelvic-Hip Region.* 3rd ed. London, England: Churchill Livingstone; 2004.
- Henderson L. Diagnosis, treatment, and lifestyle changes of interstitial cystitis. *AORN J.* 2000;71(3):525–538.
- Rovner E, Probert K, Brensinger C, et al. Treatments used in women with interstitial cystitis: the Interstitial Cystitis Data Base (ICDB) study experience. *Urology.* 2000;56(6):940–945.
- US Department of Health and Human Services. *National Kidney and Urologic Diseases Information Clearinghouse.* Bethesda, MD: National Institutes of Health; 2011.
- Ottm D, Carr L, Perks A. Interstitial cystitis and female sexual dysfunction. *Urology.* 2007;69(4):608–610.
- Nickel JC, Moldwin R, Lee S, Davis EL, Henry RA, Wyllie MG. Intravesical alkalized lidocaine (PSD597) offers sustained relief from symptoms of interstitial cystitis and painful bladder syndrome. *BJU Int.* 2009;103(7):910–918.
- Parsons L. Successful downregulation of bladder sensory nerves with combination of heparin and alkalized lidocaine in patients with interstitial cystitis. *Urology.* 2005;65(1):45–48.
- Waters M, Suleskey JF, Finkelstein LJ, Van Overbeke ME, Zizza VJ, Stommel M. Interstitial cystitis: a retrospective analysis of treatment with pentosan polysulfate and follow-up patient survey. *J Am Osteopath Assoc.* 2000;100(3S):13S–18S.
- Carrico D, Peters K. Vaginal diazepam use with urogenital pain/pelvic floor dysfunction, serum diazepam levels and efficacy data. *Urol Nurs.* 2011;31(5):279–284.
- Fitzgerald MP, Anderson RU, Potts J, et al., Randomized multicenter feasibility trial of myofascial physical therapy for treatment of urologic chronic pelvic pain syndrome. *J Urol.* 2009;182(2):570–580.
- Fitzgerald MP, Kotarinos R. Rehabilitation of the short pelvic floor, part I: background and patient evaluation. *Int Urogynecol J Pelvic Floor Dysfunct.* 2003;14(4):261–268.
- Oyama IA, Rejba A, Lukban JC, et al. Modified Thiele massage as therapeutic intervention for female patients with interstitial cystitis and high-tone pelvic floor dysfunction. *Urology.* 2004;64:862–865.
- Weiss JM. Pelvic floor myofascial trigger points: manual therapy for interstitial cystitis and the urgency-frequency syndrome. *J Urol.* 2001;166:2226–2231.



34. Holzberg A, Kellog-Spadt S, Lukban J, et al. Evaluation of transvaginal Thiele massage as a therapeutic intervention for women with interstitial cystitis. *Urology*. 2001;57(6)(suppl 1):119–120.
35. International Pelvic Pain Society. History and Physical Form for healthcare providers. [http://www.pelvicpain.org/pdf/History\\_and\\_Physical\\_Form/PPSH&PformR-MSW.pdf](http://www.pelvicpain.org/pdf/History_and_Physical_Form/PPSH&PformR-MSW.pdf). Accessed February 11, 2015.
36. Allen R, Hosker G, Smith A, et al. Pelvic floor damage and childbirth: a neurophysiological study. *Br J Obstet Gynaecol*. 1990;97:770–778.
37. Chatoor D, Emmnauel A. Constipation and evacuation disorders. *Best Pract Res Clin Gastroenterol*. 2009;23(4):517–530.
38. Harrington KL, Haskvitz EM. Managing a patient's constipation with physical therapy. *Phys Ther*. 2006;86(11):1511–1519.
39. Hanno P, Burks DA, Clemens JQ, et al. *American Urological Association Guidelines on Interstitial Cystitis/Bladder Pain Syndrome*. Linthicum, MD: American Urological Association; 2014.
40. Ware J. *User's Manual for the SF-36v2 Health Survey*. 2nd ed. Lincoln, RI: Quality Metric; 2011:81–84.
41. Melzack R. The Short-Form McGill Pain Questionnaire. *Pain*. 1987;30(2):191–197.
42. American Physical Therapy Association. Guide to Physical Therapist Practice. 2nd edition, revised. *Phys Ther*. 2003;81:9–744.
43. Haugstad GK, Haugstad TS, Kirste UM, et al. Posture, movement patterns, and body awareness in women with chronic pelvic pain. *J Psychosom Res*. 2006;61(5):637–644.
44. Jackson R. Diagnosis and treatment of pelvic girdle dysfunction. *Orthop Phys Ther Clin North Am*. 1998;7(3):413–445.
45. Magee DJ. *Orthopedic Physical Assessment* 4th ed. Philadelphia, PA: Saunders; 2002.
46. Hartman D, Sarton J. Chronic pelvic floor dysfunction. *Best Pract Res Clin Obstet Gynaecol*. 2014;28:977–990.
47. Norkin C, White DJ. *Measurement of Joint Motion: A Guide to Goniometry*. 3rd ed. Philadelphia, PA: FA Davis; 2003.
48. Kaltenborn F. *Manual Mobilization of the Joints, Volume II: The Spine*. 4th ed. Gotesburg, Sweden; 2003.
49. Karnath B. Clinical signs of low back pain. *Hosp Physician*. 2003;5:39–44.
50. Butler D, Jones M. *Mobilisation of the Nervous System*. Elsevier Health Sciences: London, England; 1991:265.
51. Lee D. The One-Leg Standing Test and the Active Straight Leg Raise Test: a clinical interpretation of two tests of load transfer through the pelvic girdle. *Orthop Div Rev*. 2005. <http://www.kalindra.com/LoadTransferTests.pdf>. Accessed February 11, 2015.
52. Hislop H, Montgomery J. *Daniels and Worthingham's Muscle Testing: Techniques of Manual Examination*. St. Louis, MO: 2002.
53. Messelink B, Benson T, Berghmans B, et al. Standardization of terminology of pelvic floor muscle function and dysfunction: report from the Pelvic Floor Clinical Assessment Group of the International Continence Society. *Neurourol Urodyn*. 2005;24:374–380.
54. Voorham-van der Zalm PJ, Lycklama A, Nijeholt GA, Elzevier HW, Putter H, Pelger RC. Diagnostic investigation of the pelvic floor: a helpful tool in the approach in patients with complaints of micturition, defecation, and/or sexual dysfunction. *J Sex Med*. 2008;5(4):864–871.
55. Goldstein A. Cotton Swab Test. <http://sexualmed.org/index.cfm/diagnostic-tests-for-women/cotton-swab-test>. Accessed February 11, 2015.
56. Goldstein I, Goldstein AT. Vestibulodynia: diagnosis. <http://www.sexualmed.org/index.cfm/sexual-health-issues-for-women/vestibulodynia/diagnosis/index.cfm>. Accessed February 11, 2015.
57. Hides JA, Stokes MJ, Saide M, Jull GA, Cooper DH. Evidence of lumbar multifidus muscles wasting ipsilateral to symptoms in patients with acute/subacute low back pain. *Spine*. 1994;19(2):165–177.
58. Hides JA, Richardson CA, Jull GA. Multifidus recovery is not automatic following resolution of acute first episode low back pain. *Spine*. 1996;21(23):2763–2769.
59. Chaitow L. Breathing pattern disorders, motor control and LBP. *J Osteopath Med*. 2004;7(1):34–41.
60. Chaitow L. Breathing pattern disorders and back pain. In: Vleeming A, Mooney V, Stoekart R, eds. *Movement Stability & Lumbopelvic Pain*. Edinburgh, Scotland: Churchill Livingstone/Elsevier; 2007:563–572.
61. Hodges P. Postural and respiratory functions of the pelvic floor muscles. *Neurourol Urodyn*. 2007;26(3):362–371.
62. Friedlander JI, Shorter B, Moldwin RM. Diet and its role in interstitial cystitis/bladder pain syndrome (IC/BPS) and comorbid conditions. *BJU Int*. 2012;109(11):1584–1591.
63. Chaiken DC, Blaivas JG, Blaivas ST. Behavioral therapy for the treatment of refractory interstitial cystitis. *J Urol*. 1993;149(6):1445–1448.
64. Anderson R, Wise D, Sawyer T, Nathanson BH. Safety and effectiveness of an internal pelvic myofascial trigger point wand for urologic chronic pelvic pain syndrome. *Clin J Pain*. 2011;27(9):764–768.